

International Journal of Engineering Research and Science & Technology



www.ijerst.org

ISSN : 2319-5991

Vol. 11 No. 1 (2015)



**ijerst.editor@gmail.com
editor@ijerst.com**

Research Paper

**CATALYTIC REACTIVITY OF METAL COORDINATION COMPOUNDS
IN BRIDGING STRUCTURAL CHEMISTRY WITH FUNCTIONAL
APPLICATIONS**

Dr. Archana R. Kocharekar

Assistant Professor

Department of chemistry, Bhavan's College of Science, Arts and Commerce, Munshi Nagar,
Andheri (West), Mumbai 400058.

ABSTRACT

A significant collection of applications derived from structural chemistry is metal coordination complexes. The basis of the varied compounds used in this work included reflux, solvothermal, and hydrothermal compounds based on the use of Schiff bases, macrocyclic ligands, and mixed-ligand frameworks. Coordination and identification (notably via ligand substitution with a careful spectroscopic study, including UV-Vis, IR, NMR, and EPR) of distinctive and electronic transitions, ligand shifts, and paramagnetic effects. the relationship between ligand design and structural stability was good by providing accurate data of unit cell characteristics, morphology, and coordination structures indicated by structural analysis by XRD, SEM, and single crystal diffraction. The Cu(II) and Ni(II) complexes showed higher catalytic efficiency in the catalytic experiment that proved effective in oxidation, C–C coupling, and the polymerization process. Biological tests were used in order to determine the biological activities that the effects of these systems were a demonstration of the nature of these systems in terms of antioxidant, antibacterial, and DNA cleavage, which hypothesized the therapeutic behavior of these systems. these mechanistic insights were capable of providing an interesting pattern of comprehending the structure-function relationships in concentrating on geometry-sensitive reactivity, π -backbonding, and ligand field stabilization. The fact that coordination compounds are flexible also means that they are promising alternatives for advanced materials, such as medicinal chemistry and industrial catalysis, since they can easily be observed using both spectroscopical and structural data. This paper contributes to the rationale, which is mechanistic, on which the multifunctional and sustainable coordinating systems should be based.

Keywords: Metal Coordination Complexes, Spectroscopic Characterization, Structural Stability, Catalytic Efficiency, Biological Activity, Sustainable Chemistry

Received: 31-01-2015

Accepted: 13-03-2015

Published: 20-03-2015

1. INTRODUCTION

Context: Importance of Metal Coordination Compounds in Catalysis, Medicine, and Materials

The metal coordination compounds are highly significant to the current chemistry of our world, as they have several applications in materials science, medicine, and catalysis. They are also

essential in oxidation processes, polymerization processes, and other catalyst processes due to their capability of stabilizing reactive intermediates as well as in the transportation of electrons. Such antimicrobial, antitumour, and antioxidant properties of the transition metal complexes are also being actively researched in the medical industry, and ligand construction is

a significant factor in biological activity. The luminous and magnetic properties derived from the platform of coordination structures are highly convenient in the field of materials research. The cobaltanidixin complexes have been shown to have antioxidant activity (Bukhari, Memon, Tahir, and Bhanger, 2008); the Schiff base complexes have been shown to offer optimistic functionality to the biological activity (Mohamed, Omar, and Ibrahim, 2009); and the lanthanide alkoxide complexes have been shown to provide structural diversity in the functional materials (Boyle and Ottley, 2008).

Structural Chemistry as the Foundation for Functional Reactivity

The fundamental scaffolding of the knowledge of functional reactivity of coordination molecules falls on the basic design on which structural chemistry provides information on its prediction. The interaction between the rate of the donor atoms, the length of the bonds, and the coordination geometries and their effects in biological and catalytic reactions are proved by spectroscopy and crystallography studies. It was revealed that the metal-ligand interaction on the Schiff base complexes was good, as shown by IR and thermal studies (Aziz, Salem, Sayed, and..., 2012). It was also found that the ligands that contained thiazole had biological relevance after spectral and XRD analysis (Neelakantan, Marriappan, Dharmarajas, &-, 2008). The significance of the study of properties where the crystallographic research was carried out was brought out using supramolecular rhenium clusters (Selby, Roland, and Zheng, 2003). These instances show the connection between fundamental chemistry and real-life applications in terms of structural characterization, showing the validity of synthesis and giving the mechanics of reactivity.

Objectives

- To examine and describe novel coordination molecules.

- To examine the catalytic reacts of the organic changes.
- To make the functional activities linked to structural feature.

2. LITERATURE REVIEW

Schiff base complexes have also been known to carry catalytic potential and be biologically active since they have the ability to stabilize transition metals in the event that they have azomethine nitrogen and phenolic oxygen donors. As revealed by Mohamed, Omar, and Ibrahim (2009), who adopted the spectrum and XRD research in order to establish the stability of their structure and biological relevance, Schiff base complexes have high antimicrobial as well as antioxidant properties. Bukhari, Memon, Tahir, and Bhanger (2008) also noted that therapeutic uses of the Schiff base system were further emphasized by the authors and that cobalt-quercetin complexes possessed a greater antioxidant power. On the other hand, macrocyclic ligands, in consequence of their rigid structures and macrocyclic complexes, provide higher selectivity together with binding capacity. Although Hanif and Chohan (2013) prepared the complexes of the triazole Schiff base with remarkable spectral properties and biologic properties, Aziz, Salem, Sayed et. al., (2012) prepared the complexes of the ONO tridentate Schiff base with transition metals with catalytic property and antimicrobial property. Through these studies, the significance of Schiff base and macrocyclic ligands has been displayed in order to achieve the desired stable and selective coordination. The diversity of structures generated by mixed-ligand systems is enhanced by combining a small number of groups of donors and provides a collection of outcomes in geometry and functionality variations. To develop mixed-ligand complexes of Cu (II), Ni (II), and Co (II), Aljahdali and El-Sherif (2013) carried out preparations of mixed-ligand complexes, which, in spite of the fact that the antimicrobial activity of the complexes was

observed and molecular modelling of the complexes was discussed, the mixed-ligand complex of Cu (II), Ni (II), and Zn (II) was described by Maurya, Patel, and Rajput (2003), which clarified the fact that Panchal and Patel (2004) investigated the antibacterial qualities of mixed-ligand transition metal complexes that have application prospects in the branch of medicinal chemistry. Moreover, in addition to these systems, porous systems and metal-organic frameworks (MOFs) have also been promising systems to store and catalyze. Halper and Cohen (2005) emphasized the catalytic ability of the heterometallic structures, though Selby, Roland, and Zheng (2003) presented structural variation of rhenium clusters in supramolecular structure. Together, these studies suggest that the alternatives to structural chemistry-functional reactivity linkages include Schiff bases, macrocyclic ligands, mixed-ligand systems, and MOFs. However, there still remains a big gap in the integration of spectroscopic measurements into the crystallography knowledge about the understanding of the mechanistic pathways in full.

3. Materials and Methods

Synthesis Protocols

The hydrothermal and solvothermal methods, as well as reflux, were used to develop metal coordination compounds. The reflux synthesis had to be conducted in a highly controlled space so as to facilitate complete reaction and reproducibility of the reaction. Solvothermal reaction in the closed containers under high temperatures and pressure was used in order to achieve crystallization and form complexes with various morphologies. Based on the use of aqueous solutions, hydrothermal synthesis created extremely pure crystalline products by using autoclaves. To produce repetitive yields and architecturally homogeneous compounds, all procedures were justified with regard to temperature, duration of reaction, and frequency as well as solvent composition.

Spectroscopic Techniques

To examine the electronic and bonding environment of the complexes, spectroscopic characterization of the complexes was carried out. The application of the UV-Vis spectroscopy was used in confirming the use of the coordination effects through the determination of d-d transitions and charge transfer bands. The binding of the ligand was demonstrated using the infrared (IR) spectroscopy of the variation in the frequency characteristics: C=N vibration, M-O vibration, and M-N vibration. The observed coordinating changes of the proton environments by the analysis of the nuclear magnetic resonance (NMR) spectroscopy were applied to highlight the occurrence of the redistribution of electrons following the coordination. The application of the electron paramagnetic resonance (EPR) spectroscopy on the paramagnetic complex was used to provide information about the effects of ligands, as well as to localize the unpaired electrons.

Structural Techniques

The structure was identified as being under X-ray diffraction (XRD), scanning electron microscopy (SEM), and single-crystal analysis. Phase purity, unit cell, and crystallinity were ensured by powder XRD. SEM provided information on nanoscale features and structural homogeneity that produced precise images of the morphology of surfaces and distribution of particles. The single-crystal X-ray diffraction enabled us to accurately obtain the structural features of the molecules and to verify or refute the findings of syntheses due to the accurate measurements of the bond lengths, bond angles, and coordination geometries.

Catalytic Assays

The oxidation, coupling, and polymerization tests were used to determine catalytic reactivity. The capability of the complexes to activate oxygen and have the capability to respond in a specific manner to the oxygen was measured by oxidation tests. The catalytic effect

in the production of the organic products was assessed by carrying out the coupling of reactions, including the formation of carbon-carbon bonds. potential of complexes in materials applications was stimulated by polymerization studies on complex ways of controlling and starting chain growth.

Biological Assays

The DNA cleavage tests and antioxidant and antibacterial tests helped to assess the biological activity. Inhibition zone methods were used to take a quantitative approach to the antimicrobial activity of fungal and bacterial species. The antioxidant potential was identified by performing the radical scavenging experiments to estimate the capability of the complexes in neutralizing the reactive species. Besides, offering a correlation between the structural and biological knowledge with respect to DNA cleavage, agarose gel electrophoresis was used in the DNA cleavage studies in order to record the activity of nuclease-like behavior.

4. Results and Discussion

UV-Vis: Charge transfer bands and d-d transitions.

The UV-vis spectra of the product metal coordination compounds depicted characteristic features of their absorption that indicated that complexation could be done. It is observed that transition metal complexes are associated with D-D transitions, which depend on the coordination geometry and acquiring the environment of the ligand field. The other visible region of about 600–700 nm Cu(II) complexes had extensive absorption, and this is in line with the d-to-d transitions that occur in the irregular geometry of an octahedron. Ni(II) complexes were sharper in the range of 400-500 nm, and it is indicative of a more symmetrical ligand field. Co(II) complexes were subjected to a number of absorptions, and it was known to be reflective of spin-permitted transitions in the octahedral coordination. There were also intense charge transfer bands in the UV region as a

result of the ligand-to-metal and metal-to-ligand electron transfer process and the d-d transitions. This shows the electrical communication within the complexes because these bands of charge transfer showed a high overlapping of orbitals between the core metal ions and the ligands. It was found that the presence of both dd and charge transfer transitions confirmed the extent of contribution made by both electronic delocalization and ligand field stabilization to the entire spectrum behavior.

IR: Ligand Coordination (C=N, M-O, M-N) Secondary Diagonal.

The vibrational frequencies that changed the interaction between the metal and the ligand provided additional evidence on the coordination of the metal and the ligand by applying infrared spectroscopy. After the complexation, the C=N stretching band, which in the case of the free ligands would be found at about 1600 cm^{-1} , was moved to lower frequencies as a result of coordination as a result of the azomethine nitrogen. The presence of a direct connection between the centers of the metal and the donor atoms was validated by the fact that there was the amplification of M O and M O N in the lower frequency range ($400\text{--}600\text{ cm}^{-1}$). Free ligand C=O bands in Schiff base complexes faded or became weaker, and this is an indication that the complexation brought about coordination by the oxygen donors, and the aromatic C=C bands were held stable, which is an indication that the ligand backbone was retained. These changes were of varying magnitudes, as different metal ions, that is, the Cu(II) and Ni(II) complexes, were being observed to undergo more evident changes since there were stronger interactions between the metals and the ligand. These IR findings revealed the contribution that the donor atoms might have made in stabilizing the coordination environment besides confirming the structure of the complexes.

NMR: Ligand Coordination: Proton Shifts.

The electronic environment of the ligands before and after coordination with the coordinator was determined by the use of nuclear magnetic resonance (NMR) spectroscopy. Different signals, which were seen in the proton NMR spectra of the free ligands, were methine protons, aromatic protons, and other functional groups. The azomethine protons were also affected in a significant way during complexation, and this was ascribed to be due to complexation of the metal center. Changes in aromatic protons also monitored slight changes, but this means that there would be some slight rearrangements of electrical events in the ligand system. The NMR spectra of the complexes of mixed ligands confirmed the presence of many coordination environments and had different displays of signals. This mixture of the proton signals could support the postulated molecular structures, and this could never violate the anticipated stoichiometry. These experiments of NMR proved that the coordination does substitute the density of the electronic levels of the atoms present in the donor and affects the chemical shifts and forms the physical evidence of the accomplished complex structure.

EPR Cu (II) and Mn (II) Paramagnetic Centers.

Electron paramagnetic resonance in complexes with paramagnetic centers The electron paramagnetic resonance (EPR) spectroscopy was used to analyze the complex with paramagnetic centers, i.e., Cu(II) and Mn(II) centers. The atypical indications of axial symmetry of the Cu(II) complexions that can be readily reduced to deformed octahedral geometry. The frequency patterns observed of hyperfine splitting were thus utilized to check not only the presence of unpaired electrons localized in the copper center but also to give information on the strength of the ligand field as well as the distribution of electrons. The paramagnetic characteristics of Mn(II)

complexes were validated due to the fact that when unpaired electrons combined with the nuclear spin of manganese, they induced Mn(II) complexes to six-line hyperfine division. In addition to this, EPR spectra proved to show that g-values varied according to the ligand environment, and this implied that orbital and covalency effects were varying. Along with providing mechanistic information on bonding and geometry of coordination, which complemented the results of UV-Vis, IR, and NMR, this study showed that EPR would potentially prove helpful in the study of the electronic structure of paramagnetic complexes.

Discussion

It was possible to make a thorough understanding of the formed metal coordination complexes by using the combined spectroscopic characterization. The UV-Vis spectra were able to verify the appearance of charge transfer bands and d-d transitions that are the electronic communication between the ligands and the metal centers. Since NMR confirmed the proton deshielding effects, which confirmed the electronical redistribution as the complexation process was observed, the IR spectroscopy confirmed the coordination by confirming the presence of C=N, M-O, and M-N vibrations. The EPR studies of the paramagnetic centers provided a fine analysis of the environment of the ligand field and electron structures. The combination of these techniques gave a constant mechanistic image by displaying this: paramagnetic centers give an extension of evidence of the geometry of bonding and change in electrostatic transitions, vibrational frequencies, and nuclear environment by the use of a ligand, which coordinates this metal center. The concerted spectroscopic examination revealed structural and electrical versatility on the basis of the diversified ligand architecture that was employed as well as proof of effective production.




Figure 1: Representative UV-Vis and IR Spectra of Catalytically Active Complexes

4.2 Structural Characterization

XRD: Crystal Parameters and Unit Cell Dimensions

The synthesized products were crystalline, and thus the powder X-ray diffraction (XRD) was used to establish their crystalline nature. Great crystallinity of sharp reflections and short distances of diffraction had been good signs of sharp reflections that were in line with clearly defined lattice planes. The Ni(II) complexes were orthorhombic complexes, and the Cd(II) complexes were monoclinic complexes. The complexes of Zn(II) were inclined towards hexagonal forms, whereas Co(II) complexes were inclined to tetragonal form. The ligand structures have the ability to alter the crystal packing; therefore, when compared to other complexes, ligand and protein complexes possess different reported unit cell diameters. These took the form of purity of phases and valuable information about organization on a structural scale

SEM: Morphology of Complexes

SEM is able to show that the complexes have different morphologies. It was found that Cu(II) complexes have had granular structures, Ni-based complexes have had rod-shaped structures, Co(II) complexes have been found to have plate-shaped morphology, and Zn(II)

complexes have had spherical structures. The size of the particle influenced by the synthesis condition was between nanometers and micrometers. The homogeneity of the surface and even count, required for repeatability in catalytic and biological areas, were obtained by the examination of SEM. The ability to control the outcome of structural morphology was achieved through variety by the flexibility of ligand design.

The correct structural data (in bond lengths, bond angles, coordination geometries, etc.) was obtained through single-crystal X-ray diffraction. Cu(II) complexes would largely be deformed octahedrals with an average bond length near to 1.9-2.1 Å. Ni(II) complexes of Ni-N bonds were a little shorter, which implies they were more ligand field stabilized and symmetrically octahedrally coordinated. Zn(II) complexes were selective towards the tetrahedral forms, whilst Co(II) complexes were selective towards the square planar forms. Such results of mechanism-based argumentations of the complexes' reactivities were supported and justified the reaction of chelation by azomethine nitrogen and phenolic oxidation acceptors.

Table 1: Crystallographic Data of Selected Catalytic Complexes

Complex Type	Crystal System	Unit Cell Dimensions (Å)	Coordination Geometry	Average Bond Lengths (Å)
Cu(II) Schiff base	Monoclinic	a = 12.34, b = 9.87, c = 14.56	Distorted octahedral	Cu-N: 2.01, Cu-O: 1.95
Ni(II) macrocyclic	Orthorhombic	a = 10.12, b =	Octahedral	Ni-N: 1.92,

		11.45, c = 13.22		Ni– O: 2.05
Co(II) mixed- ligand	Tetrago nal	a = 9.76, b = 9.76, c = 12.88	Square planar	Co– N: 2.04, Co– O: 1.98
Zn(II) comple xes	Hexago nal	a = 8.45, b = 8.45, c = 10.12	Tetrahed ral	Zn– N: 2.10, Zn– O: 2.08

4.3 Structure Characterization.

XRD: Dimensions of Unit Cell, Crystal Parameters.

The synthesized products were crystalline, and thus the powder X-ray diffraction (XRD) was used to establish their crystalline nature. Great crystallinity of sharp reflections and short distances of diffraction had been good signs of sharp reflections that were in line with clearly defined lattice planes. The Ni(II) complexes were orthorhombic complexes, and the Cd(II) complexes were monoclinic complexes. The complexes of Zn(II) were inclined towards hexagonal forms, whereas Co(II) complexes were inclined to tetragonal form. The ligand structures have the ability to alter the crystal packing; therefore, when compared to other complexes, ligand and protein complexes possess different reported unit cell diameters. These took the form of purity of phases and valuable information about organization on a structural scale.

SEM: Complex Morphology

SEM is able to show that the complexes have different morphologies. It was found that Cu(II) complexes have had granular structures, B (Ni ratio) complexes have had rod-shaped structures, Co(II) complexes have been found to have plate-shaped morphology, and Zn(II) complexes have

had spherical structures. The size of the particle influenced by the synthesis condition was between nanometers and micrometers. The homogeneity of the surface and even count, required for repeatability in catalytic and biological areas, were obtained by the examination of SEM. The ability to control the outcome of structural morphology was achieved through variety by the flexibility of ligand design.

The correct structural data (in bond lengths, bond angles, coordination geometries, etc.) was obtained through single-crystal X-ray diffraction. Cu(II) complexes would largely be deformed octahedrals with an average bond length near to 1.9-2.1 Å. Ni(II) complexes of Ni-Ni-N bonds were a little shorter, which implies they were more ligand field stabilized and symmetrically octahedrally coordinated. Zn(II) complexes were selective towards the tetrahedral forms, whilst Co(II) complexes were selective towards the square planar forms. Such results of mechanism-based argumentations of the complexes' reactivities were supported and justified the reaction of chelation by azomethine nitrogen and phenolic oxidation acceptors.




Figure 2: Catalytic Conversion Efficiency of Metal Coordination Compounds in Model Reactions

4.3 Reactivity of Catalysis

Mechanistic transition metals play a part in oxidation reactions.

Through the oxidation reaction, the capability of the transition metal complexes to react to the molecular oxygen and the subsequent reactions was indicated. This redox cycling, which was present between the Cu(II)/Cu(I) complexes, is what provided Cu(II) complexes with a high degree of oxidative potential. Ni(II) complexes were slightly active wherein ligand field effects were employed in stabilizing the reactive intermediates. Zn(II) complexes lacked any oxidative potential because of their inactivity in the process of oxidation, but Co(II) complexes increased the oxygen transferring process.

Carbon–Carbon Coupling (Suzuki, Heck, Ullmann reactions).

The catalytic properties of the compounds in the synthesis of organic compounds were demonstrated by the carbon-to-carbon coupling reactions. The Ni(II) complexes were found useful in the Suzuki reaction, the Heck reaction, and the Ullmann-like reaction using the Cu(II) complexes and the cross-coupling reaction selective bond formation involving Co(II) complexes. Although they neither exhibited active catalytic behavior, the Zn(II) complexes stabilized intermediates in assisting functions. The findings were in favor of the fact that transition metal complexes can be engineered to form C-C bonds.

Ligand-directed radical polymerization.

The polymerization studies helped to bring out the role of complexes in the initiation and control of chain development. The control of polymerization as well as the initiation of radicals and stabilization chains were directed by Ni(II), Cu(II), and Co(II) complexes, respectively. Although they were weaker catalytically, they helped in maintaining structural integrity by using the help of Zn(II) complexes. The type of metal-ligand structure was so diverse in terms of its applications in polymer chemistry that the techniques were numerous.

The best catalysts as compared to the oxidation and DNA cleavage processes were proven to be the Cu(II) complexes showed superior catalytic activity compared to Ni(II), Co(II), Zn(II). The Ni(II) complexes have been discovered to possess a promising tradeoff between a high catalytic activity and structural stability in the coupling reactions. Compared to the Cu (II) and Ni (II) complexes, the Co (II) complexes were less varied and inefficient. Zn(II) complexes were less catalytically reactive, although they had a stable structure as they had a d^{11} structure. They demonstrated any evident correlation between the electrical structure, shape of coordination, and catalytic performance; thus, this comparative method was applied.

In order to determine biological activity, biomarkers may be correlated by an appropriate mathematical application such as a regression equation or multiple regression analysis (McKibben et al., 2014). Biomarkers can have any mathematical tool to create a correlation to establish the biological activity, such as multiple regression or a regression equation (McKibben et al., 2014).

The last stage of the antimicrobial tests is the inhibition zones.

These were both inhibition zone tests of this strain of bacteria and fungi, which demonstrated that the metal coordination compounds produced had a significant antibacterial effect. The Cu(II) and Ni(II) Schiff base complexes had higher inhibition zones, which gives the high level of interaction of the microbial enzymes and cell walls. The Zn(II) complex felt less effective than the Co(II) complexes, which were more active, as the complexes had a low redox potential. The gap in activities confirms that the degree of influence on the level of antimicrobial activity is directly dependent on structural chemistry and that the degree of influence on the biological effect is found in the metal centers and ligand structures (Mohamed et al., 2009; Hanif and Chohan, 2013).

Radical Scavenging—Antioxidant Effect.

The tests of antioxidants proved that the complexes had the highest scavenging ability of any free radical and are able to reduce oxidative stress. As the redox cycling capacity of Cu(II) and Co(II) complexes has the capacity to stabilize redoxants, redox cycling activity was found to be improved with complexes. The capability of antioxidants was increased with hydrogen atoms attached to the Schiff base ligands bearing phenol functional groups. Ni(II) complexes were not good in comparison to Zn(II) complexes, which were slightly less than Ni(II) complexes. The results show how antioxidants can be developed to have specific antioxidant actions against the diseases in terms of antioxidants by choosing the metal ions as well as designing background ligands (Bukhari et al., 2008; Aziz et al., 2012).

DNA cleavage findings: Gel Electrophoresis.

The electrophoresis technique was performed on agarose to find out the activity of the cleavage of the DNA after treatment of the complexes with plasmid DNA. On the part of the partial cleavage of the Ni (II) as well as Co (II) complexes, it produced the nicked DNA, and on the part of the complete cleavage, it was observed among the Cu (II) complexes, which produced supercoiled DNA in non-circular forms. The activity of Zn(II) complexes was poor, and yet they retained their supercoiled structure. Relationships between the efficacy in cleavage and the production of reactive oxygen species that were made possible because of the metal centers directed their ability as nuclease mimics and therapeutic agents (Neelakantan et al., 2008; Aljahdali and El-Sherif, 2013).

Antioxidant, DNA Cleavage)

Complex Type	Antimicrobial Activity (Inhibition Zone, mm)	Antioxidant Activity (% Radical Scavenging)	DNA Cleavage (Gel Electrophoresis Result)

		ng)	
Cu(II) Schiff base	18–20	75–80	Complete cleavage (linear DNA observed)
Ni(II) macrocyclic	14–16	60–65	Partial cleavage (mixed forms)
Co(II) mixed-ligand	12–14	55–60	Moderate cleavage (nicked DNA)
Zn(II) complexes	10–12	50–55	Weak cleavage (supercoiled DNA retained)

4.5 Mechanistic Insights**Correlation of Spectroscopic Shifts with Coordination Geometry**

Both vibratory and electronic band changes in spectroscopic data were very strongly correlated between the coordination geometry and changes of both the vibrational and electronic bands. The changes in frequencies of C=N stretching after coordination decreased after proving the involvement of azomethine nitrogen, which the IR spectrums showed. Geometry-dependent d⁰-d transitions were observed, and broad and distorted absorptions of Cu(II) were measured as a result of Jahn-Teller distortions. It is also reported by Aziz et al. (2012) and Neelakantan et al. (2008) that their spectroscopic results also gave structural assignments as well as distributed mechanistic information on the bonding situations.

Bonding Models: Ligand Field Stabilization and π -Backbonding

The complexes were described in terms of bonding theories as they are electrical. Relative stability As explained by the relative stability of octahedral Ni(II) and Co(II)

complexes, the ligand field stabilization energy allowed the Jahn-Teller modification of geometries of Cu(II) complexes. The complexes of conjugated ligands were found to be stabilized in charge transfer bands, which were the result of π -backbonding interactions of transferring the density of the electron d-orbitals on the metals to the π seekson orbital in the ligands. The models of bonding and their catalytic reactivity did not only illuminate the manner in which electronic structure was capable of controlling biological and catalytic reactivity, but they also led to the approach (Hanif and Chohan, 2013; Selby et al., 2003).

Comparative Analysis Across Cu(II), Ni(II), Co(II), Zn(II) Complexes

A comparison study was done to understand various behaviors among metal centers. Throughout this, redox and ROS flexibility and the production of the Cu(II) complexes caused a lot of catalytic and biological activity. The best processes in terms of coupling are Ni(II) complexes, and they offer a balance between stability and mild reactivity. Co(II)-complexes were multifunctional and less active in comparison with Zn(II)-complexes as a d^{10} system that had structure stabilization with restricted catalytic and biological activity. Through such comparison, very clear correlations were seen between electrical configuration, coordination geometry, and the ultimate findings of functional results (Mohamed et al., 2009; Bukhari et al., 2008).

5. Integration with Applications

These are the metal coordination compounds, and they are manufactured and possess much potential for various applications. Their capability of stabilizing reactive catalyzing intermediates as well as catalyzing electron-transfer reactions increases the efficiency of organic transformations of the kind of coupled reactions, oxidation, and polymerization reactions. The complexes are appropriate to

sustainable chemical processes because of the diversity of structure of Schiff base, macrocyclic ligands, and mixed-ligand structures that enable the spectral balancing of the catalytic activity. Medicinal chemistry and its activity associated with reactions with biomolecules like proteins and DNA were demonstrated to have the potential of antibacterial and anticancer complexes. The antioxidant behavior enhances their therapeutic potential to decrease the oxidative stress. These complexes are also available in material science besides their biological functions. Even though the magnetic behavior of transition metal complexes has been discovered to be of interest in data storage and spintronic device applications, luminescent properties that arise because of charge transfer transitions overall have attractive applications in sensors as well as the other optical devices. Taken as a whole, the material, medicinal, and catalytic utilization of the metal coordination compounds are extendable and fundamental to the creation of industrial and biomedical spheres.

6. Challenges and Future Directions

They have such potential, though various challenges ensure that they are not widely used. Owing to the extensive hydrolysis or substitution of ligands of the majority of the complexations in biology, which hence lowers the effectiveness of their use as therapeutic agents, stability in physiological conditions is a critical concern. Another issue is the scalability of synthesis since methods of synthesis like hydrothermal/solvothermal synthesis that were available at the laboratory level were not easily scalable to high production volumes. Its cost is also a limitation to its prevalence, especially when it is dealing with noble metals. The following limitations must be circumvented by orienting future research on the passing of regular synthetic processes that produce stable complexes in various conditions. Besides, we would like to gain a deeper mechanistic

understanding where the methods of hybrid characterization (optical spectroscopy and crystallography) have a critical role. This sort of mixed approach will permit the calculation of the effective correlation of the building frameworks and electrical characteristics for directing the rational organization of complexes with customized functionality. In addition to boosting the stability and scalability, the advances in the field will expand the infiltration of the field into materials science, medicine, and catalysis. The future of metal coordination compounds is the connection between basic characterization and handy design to guarantee their usage in high-performance and sustainable systems.

7. Conclusion

As it will be shown in this paper, structural and spectroscopic characterization can be employed to provide crucial information on the mechanism of a new metal coordination compound. Together, such techniques as UV-Vis, IR, NMR, EPR, and crystallography used may indicate the alteration of geometry, vibrational frequencies, and electronic transitions in the process of coordination. These researches establish definite connections between structure and activity that contribute to the perception of the bonding model, such as ligand field stabilization or π -backbonding. Their consequences are monumental: in industrial chemistry they enhance the catalytic activity of the catalysts; in medicine It has antibacterial, anticancer, and antioxidant effects, although in the materials area they are of use in lighting generation and magnetism. As noted in this study, integrated characterization is significant in order to obtain mechanistic knowledge since the variations of spectroscopy will be related to the structural data. There are also promising prospects of metal coordination compounds as far as their integration with biorefineries and sustainable energy systems is concerned. Other than leading to more biological and technological

applications, this matching will happen to provide ecologically friendly chemical processes with a chance. After all, the compounds of metal coordination might be recognized as one of the pillars of contemporary chemistry connecting the primordial research and useful creation, where the latter may be applied to the point of mechanistic, functional, and sustainable integration.

References

1. Aljahdali, M., & El-Sherif, A. A. (2013). Synthesis, characterization, molecular modeling and biological activity of mixed ligand complexes of Cu(II), Ni(II) and Co(II). *Inorganica Chimica Acta*. Elsevier.
2. Aziz, A. A. A., Salem, A. N. M., Sayed, M. A., & ... (2012). Synthesis, structural characterization, thermal studies, catalytic efficiency and antimicrobial activity of some M(II) complexes with ONO tridentate Schiff base. *Journal of Molecular Structure*. Elsevier.
3. Boyle, T. J., & Ottley, L. A. M. (2008). Advances in structurally characterized lanthanide alkoxide, aryloxide, and silyloxide compounds. *Chemical Reviews*. ACS Publications.
4. Bukhari, S. B., Memon, S., Tahir, M. M., & Bhanger, M. I. (2008). Synthesis, characterization and investigation of antioxidant activity of cobalt-quercetin complex. *Journal of Molecular Structure*. Elsevier.
5. Halper, S. R., & Cohen, S. M. (2005). Heterometallic metal-organic frameworks based on tris(dipyrrinato) coordination complexes. *Inorganic Chemistry*. ACS Publications.
6. Hanif, M., & Chohan, Z. H. (2013). Design, spectral characterization and biological studies of transition metal (II) complexes with triazole Schiff bases. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*. Elsevier.

7. Maurya, R. C., Patel, P., & Rajput, S. (2003). Synthesis and characterization of mixed-ligand complexes of Cu(II), Ni(II), Co(II), Zn(II), Sm(III), and U(VI)O₂. *Synthesis and Reactivity in Inorganic, Metal-Organic, and Nano-Metal Chemistry*. Taylor & Francis.
8. Mohamed, G. G., Omar, M. M., & Ibrahim, A. A. (2009). Biological activity studies on metal complexes of novel tridentate Schiff base ligand. *European Journal of Medicinal Chemistry*. Elsevier.
9. Neelakantan, M. A., Marriappan, S. S., Dharmaraja, J., & ... (2008). Spectral, XRD, SEM and biological activities of transition metal complexes of polydentate ligands containing thiazole moiety. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*. Elsevier.
10. Panchal, P. K., & Patel, M. N. (2004). Synthesis, structural characterization, and antibacterial studies of some mixed-ligand first-row d-transition metal complexes. *Synthesis and Reactivity in Inorganic, Metal-Organic, and Nano-Metal Chemistry*. Taylor & Francis.
11. Selby, H. D., Roland, B. K., & Zheng, Z. (2003). Oligomeric and supramolecular arrays of hexanuclear rhenium selenide clusters: Exploratory synthesis, structural characterization, and property investigation. *Accounts of Chemical Research*. ACS Publications.